



Darganfod Hen Dai Cymreig Discovering Old Welsh Houses

Reports: Welcome to the many Discovering Old Welsh Houses Reports which are available here on our website. All the reports - House Histories, Building Reports and Tree-ring Dating reports - can be accessed - [here](#)

Discovering Old Welsh Houses studies and celebrates the traditional houses of North Wales and the lives of the people who lived in them.

The copyright of most of these reports belongs to Discovering Old Welsh Houses. Where copyright resides with others, we have made every effort to obtain their permission to reproduce reports on our site. Our policy is to allow free access to our research documents as part of the public benefit we provide as a registered charity. You are welcome to reproduce this material but if you do so, please acknowledge the source

If you find the content useful, please consider becoming a [Member](#) to access the many benefits available.



Please note that these reports are being updated as part of an ongoing programme of revision. Older reports sometimes refer to the old names of the Group. Between 2005 and 2012 also known as The Snowdonia Dendrochronology Project, then the N W Wales Dendrochronology Project and then the Dating Old Welsh Houses Group.

New reports will be added from time to time. Keep an eye on our website for updates.



©Discovering Old Welsh Houses Group

Rhif Elusen Gofrestredig: No: 1131782: Registered charity

www.discoveringoldwelshhouses.co.uk

Oxford Dendrochronology Laboratory
Report 2018/39

**DENDROCHRONOLOGICAL INVESTIGATIONS
OF TIMBERS AT PENTRE ISAF,
BRYNEGLWYS,
DENBIGHSHIRE
(SJ 1347 4726)**



Summary

Samples were taken from the east cruck to the south truss and both east and west crucks of the north truss, but were found to have very few rings. Two samples from the east cruck to the south truss matched each other and were combined to give a 52-year sequence which did not date. The other samples each had less than 40 rings and were not measured. The site, which had been previously assessed as being very marginal, remains undated.

Author: Dr M. C. Bridge FSA
Oxford Dendrochronology Laboratory
Mill Farm
Mapledurham
Oxfordshire
RG4 7TX

September 2018

Dendrochronological Investigations of timber at Pentre Isaf, Bryneglwys, Denbighshire (SJ 1347 4726)

BACKGROUND TO DENDROCHRONOLOGY

The basis of dendrochronological dating is that trees of the same species, growing at the same time, in similar habitats, produce similar ring-width patterns. These patterns of varying ring-widths are unique to the period of growth. Each tree naturally has its own pattern superimposed on the basic 'signal', resulting from genetic variations in the response to external stimuli, the changing competitive regime between trees, damage, disease, management etc.

In much of Britain the major influence on the growth of a species like oak is, however, the weather conditions experienced from season to season. By taking several contemporaneous samples from a building or other timber structure, it is often possible to cross-match the ring-width patterns, and by averaging the values for the sequences, maximise the common signal between trees. The resulting 'site chronology' may then be compared with existing 'master' or 'reference' chronologies. These include chronologies made by colleagues in other countries, most notably areas such as modern Poland, which have proved to be the source of many boards used in the construction of doors and chests, and for oil paintings before the widespread use of canvas.

This process can be done by a trained dendrochronologist using plots of the ring-widths and comparing them visually, which also serves as a check on measuring procedures. It is essentially a statistical process, and therefore requires sufficiently long sequences for one to be confident in the results. There is no defined minimum length of a tree-ring series that can be confidently cross-matched, but as a working hypothesis most dendrochronologists use series longer than at least fifty years.


The dendrochronologist also uses objective statistical comparison techniques, these having the same constraints. The statistical comparison is based on programs by Baillie & Pilcher (1973, 1984) and uses the Student's *t*-test. The *t*-test compares the actual difference between two means in relation to the variation in the data, and is an established statistical technique for looking at the significance of matching between two datasets that has been adopted by dendrochronologists. The values of '*t*' which give an acceptable match have been the subject of some debate; originally values above 3.5 being regarded as acceptable (given at least 100 years of overlapping rings) but now 4.0 is often taken as the base value in oak studies. Higher values are usually found with matching pine sequences. It is possible for a random set of numbers to give an apparently acceptable statistical match against a single reference curve – although the visual analysis of plots of the two series usually shows the trained eye the reality of this match. When a series of ring-widths gives strong statistical matches in the same position against a number of independent chronologies the series becomes dated with an extremely high level of confidence.

One can develop long reference chronologies by cross-matching the innermost rings of modern timbers with the outermost rings of older timbers successively back in time, adding data from numerous sites. Data now exist covering many thousands of years and it is, in theory, possible to match a sequence of unknown date to this reference material.

It follows from what has been stated above that the chances of matching a single sequence are not as great as for matching a tree-ring series derived from many individuals, since the process of aggregating individual series will remove variation unique to an individual tree, and reinforce the common signal resulting from widespread influences such as the weather. However, a single sequence can be successfully dated, particularly if it has a long ring sequence.

Growth characteristics vary over space and time, trees in south-eastern England generally growing comparatively quickly and with less year-to-year variation than in many other regions (Bridge, 1988). This means that even comparatively large timbers in this region often exhibit few annual rings and are less useful for dating by this technique.

When interpreting the information derived from the dating exercise it is important to take into account such factors as the presence or absence of sapwood on the sample(s), which indicates the outer margins of the tree. Where no sapwood is present it may not be possible to determine how much wood has been removed, and one can therefore only give a date after which the original tree must have been felled. Where the bark is still present on the timber, the year, and even the time of year of felling can be determined. In the case of incomplete sapwood, one can estimate the number of rings likely to have been on the timber by relating it to populations of living and historical timbers to give a statistically valid range of years within which the tree was felled. For this region the estimate used is that 95% of oaks will have a sapwood ring number in the range 11 – 41 (Miles 1997).



Section of tree with conversion methods showing three types of sapwood retention resulting in **A** *terminus post quem*, **B** a felling date range, and **C** a precise felling date. Enlarged area **D** shows the outermost rings of the sapwood with growing seasons (Miles 1997, 42)

PENTRE ISAF

A late medieval cruck-framed hall house, investigated by Peter Smith in 1976. Two cruck frames remain, with a partition in the northern truss. NPRN 27649.

SAMPLING

The timbers were investigated during August 2018, having been previously assessed as very marginal for dendro dating earlier in the year. Core samples were extracted using a 15mm diameter borer attached to an electric drill. They were labelled with the prefix **ptis**, and taken away for subsequent analysis, where they were glued to laths.

The samples were polished with progressively finer grits down to 400 to allow the measurement of ring-widths to the nearest 0.01 mm. The samples were measured under a binocular microscope on a purpose-

built moving stage with a linear transducer, attached to a desktop computer. Measurements and subsequent analysis were carried out using programs by Ian Tyers (Tyers 2004).

RESULTS AND DISCUSSION

The locations and details of the samples are described in Table 1. Two cores were taken from the east cruck to the south truss to maximise the number of rings and get sapwood. These two series were matched visually and combined into a single 52-year long sequence, but this failed to give consistent acceptable matches with the dated reference material, and remains undated. The other two cores each had less than 40 rings and were not considered useful for further analysis. The site had been assessed earlier in the year as being very marginal. Although success at a nearby site prompted this investigation.

Table 1: Details of samples taken from Petre Isaf, Bryneglwys.

Sample number	Timber and position	Sapwood complement	No of rings	Mean width (mm)	Std devn (mm)	Mean sens
Western crosswing						
ptis01a	East cruck to south truss	-	35	2.24	0.55	0.15
ptis01b	<i>ditto</i>	5	52	1.91	0.49	0.21
ptis01	Mean of 01a and 01b	5	52	2.01	0.40	0.19
ptis02	West cruck to north truss	6	<40	NM	-	-
ptis03	East cruck to north truss	-	<40	NM	-	-

Key: H/S bdry = heartwood/sapwood boundary - last heartwood ring date; std devn = standard deviation; mean sens = mean sensitivity; NM = not measured.

ACKNOWLEDGEMENTS

This report was commissioned by The Discovering Old Welsh Houses Group. I thank the members of DOWHG who made arrangements for our visits, and assisted during the fieldwork. I thank the owners (Mr and Mrs Miers), and also my fellow dendrochronologists for permission to use their data.

DOWHG wishes to acknowledge the assistance of the Woodtiger Fund, Clwydian Range and Dee Valley AONB, and the Marc Fitch Fund, towards this work.

REFERENCES

- Baillie, M.G.L. and Pilcher, J.R. (1973) *A simple cross-dating program for tree-ring research*. **Tree Ring Bulletin**, 33, 7-14.
- Bridge, M. C. (1988) *The dendrochronological dating of buildings in southern England*, **Medieval Archaeology**, 32, 166-174.
- English Heritage (1998) *Guidelines on producing and interpreting dendrochronological dates*, **English Heritage, London**.
- Miles, D. (1997) *The interpretation, presentation, and use of tree-ring dates*, **Vernacular Architecture**, 28, 40-56.
- Tyers, I. (2004) *Dendro for Windows Program Guide 3rd edn*, **ARCUS Report**, 500b.